arXiv:1301.6240v1 [math.GR] 26 Jan 2013

THE EXACT NUMBER OF r-REGULAR ELEMENTS IN FINITE EXCEPTIONAL GROUPS

SIMON GUEST

Abstract. We calculate the precise number of r-regular elements in the finite exceptional groups. As a corollary we find that the proportion of r-regular elements is at least $3577/18432$ and for all $\epsilon > 0$, there are infinitely finite simple exceptional groups such that the proportion of r-regular elements is less than $3577/18432 + \epsilon$ for some prime r.

1. Introduction

If G is a finite group, and r is a prime, then an element of G is called r-regular if its order is not divisible by r, and r-singular otherwise. Obtaining bounds (in particular lower bounds) on the number of r-regular elements in finite classical and finite exceptional groups is important for the design and implementation of various algorithms in computational group theory. In [BPS09], it is shown that the proportion of r-regular elements in a finite simple classical group is at least $1/2d$, where d is the dimension of the natural module. They show also that for every family of finite simple exceptional groups $X(q)$, there exists a constant $c(X) \geq 1/31$ such that the proportion of r-regular elements in $X(q)$ is at least $c(X)$. In [BGPW], the bounds for classical groups are improved using the quokka-set method developed in [NP10] (and inspired by [IKS95, Leh98]), and it is shown that these bounds are in some sense best possible. We use the same methods here for the finite exceptional groups. Since the Weyl group W is fixed for each family X, and the F-classes of W and corresponding maximal tori are known for each family of exceptional groups, it is possible to obtain exact results.

Theorem 1. *The precise proportions of r-regular elements in the finite exceptional groups are given in Tables 2—11.*

Corollary 1. *Let X be a family of finite simple exceptional groups $X(q)$. The proportion of r-regular elements in $X(q)$ is at least $c(X)$, where $c(X)$ is given in Table 1. Moreover, for all $\epsilon > 0$, there exists infinitely many q for which the proportion of r-regular elements in $X(q)$ is less than $c(X) + \epsilon$ for some prime r.*

It is also interesting to note that the proportion of odd order (that is, 2-regular) elements can be bounded below by a constant.

Corollary 2. *If q is odd then the proportion of odd order elements in the finite simple exceptional group $X(q)$ is at least $3577/18432$. Moreover, for all $\epsilon > 0$, there exist infinitely many $X(q)$ for which the proportion of odd order elements is less than $3577/18432 + \epsilon$.*

2. Preliminaries

In this section we give some useful technical results coming from algebraic group theory and number theory. First we set up some notation. Let $\overline{G}$ be a connected reductive algebraic group defined over $\bar{\mathbb{F}}_q$, the algebraic closure of a field $\mathbb{F}_q$ of order q. Let F be a Frobenius morphism of $\overline{G}$, and let $G = \overline{G}^F$ be the subgroup of $\overline{G}$ fixed elementwise by F, so that G is a finite group

$X(q)$	$c(X)$
${}^2G_2(q)$	$1/2$
${}^2B_2(q)$	$1/2$
${}^3D_4(q)$	$17/48$
${}^2F_4(q)$	$7/16$
$G_2(q)$	$11/36$
$F_4(q)$	$3577/18432$
$E_6(q)$	$281/1296$
${}^2E_6(q)$	$281/1296$
$E_7(q)$	$131491/589824$
$E_8(q)$	$5927482903/25480396800$

Table 1. Constants $c(X)$ for Corollary 1

of Lie type. Let $\overline{T}$ be an F-stable maximal torus in $\overline{G}$ and let $W := N_{\overline{G}}(\overline{T})/\overline{T}$ denote the Weyl group of $\overline{G}$. We will say that two elements w, w' in W are F-conjugate if there exists x in W such that $w' = x^{-1}wF(x)$. This is an equivalence relation, and we will refer to the equivalence classes as *F-classes*. Moreover, there is an explicit one-to-one correspondence between the F-classes of W and the G-conjugacy classes of maximal tori in G.

2.1. **The quokka set method.**

Now recall that every element g in G can be expressed uniquely in the form $g = su$, where $s \in G$ is semisimple, $u \in G$ is unipotent and $su = us$. This is the multiplicative Jordan decomposition of g (see [Car85, p. 11]). We now define a quokka set as a subset of a finite group of Lie type that satisfies certain closure properties.

Definition 1. Suppose that G is a finite group of Lie type. A nonempty subset Q of G is called a *quokka set*, or quokka subset of G, if the following two conditions hold.

(i) For each $g \in G$ with Jordan decomposition $g = su = us$, where s is the semisimple part of g and u the unipotent part of g, the element g is contained in Q if and only if s is contained in Q; and

(ii) the set Q is a union of G-conjugacy classes.

For G a finite group, and a prime r not dividing q, define the subset

$$Q(r, G) := \{g \in G : r \nmid |g|\},$$

consisting of all the r-regular elements g in G. We see readily that $Q(r, G)$ is a quokka set when G is a finite exceptional group. We now quote [NP10, Theorem 1.3], which will be our main tool.

Theorem 2. *Let G, W, $Q(r, G)$ be as above. For each F-class C in W, let T_C denote a maximal torus corresponding to C. Then*

$$\frac{|Q(r,G)|}{|G|} = \sum_{F-\text{classes } C \text{ in } W} \frac{|C|}{|W|}.\frac{|T_C \cap Q(r,G)|}{|T_C|}. \tag{1}$$

For an integer k and a prime r, we denote by k_r the r-part of k (that is, the largest power of r that divides k).

Lemma 1. *Let G be a finite simply connected exceptional group with centre Z. Then the proportion of r-regular elements in the finite simple exceptional group $G/Z(G)$ is $|Q(r, G)||Z|_r/|G|$.*

Proof. This is an easy consequence of [BGPW, Lemma 2.3]. □

2.2. **Cyclotomic polynomials.** Recall that the ith cyclotomic polynomial $\Phi_i(q)$ is defined to be the the unique irreducible polynomial with integer coefficients that divides $q^i - 1$ and does not a divide $q^j - 1$ for any $j < i$. The $\Phi_i(q)$ satisfy the equation

$$q^i - 1 = \prod_{j|i} \Phi_j(q). \tag{2}$$

The orders of maximal tori in the (simply connected) untwisted and Steinberg exceptional groups are products of cyclotomic polynomials (see [MT11, p. 222] for example). It is easy to see that the proportion of r-regular elements in a maximal torus T is $1/|T|_r$. In view of (**??**), the following elementary lemma on r-parts of cyclotomic polynomials will therefore be useful.

Lemma 1. *Let r be a prime and q a prime power with $(r, q) = 1$. Let e be the smallest positive integer such that $r|q^e - 1$. In particular, e is the multiplicative order of q modulo r. Let $\phi_{i,r}$ be the r-part of the ith cyclotomic polynomial $\Phi_i(q)$. If $r \geq 3$, then*

$$\phi_{i,r} = (\Phi_i(q))_r = \begin{cases} (q^e - 1)_r & \text{if } i = e; \\ r & \text{if } i = er^f \text{ and } f \geq 1; \\ 1 & \text{otherwise.} \end{cases}$$

If $r = 2$ we have

$$\phi_{i,2} = (\Phi_i(q))_2 = \begin{cases} (q-1)_2 & \text{if } i = 1; \\ (q+1)_2 & \text{if } i = 2; \\ 2 & \text{if } i = 2^f \text{ and } f \geq 2; \\ 1 & \text{otherwise.} \end{cases}$$

Remark 1. Fermat's little theorem implies that $r - 1 = ke$ for some integer k. In particular $r = ke + 1 \geq e + 1$.

Proof. This follows from (**??**) and the formula for the r-part of $q^i - 1$ given, for example, in [BGPW, Lemma 2.5]. □

3. Proof of Theorem 1

Recall that e is the multiplicative order of q modulo r. We denote by $\phi_{i,r}$ the r-part of the ith cyclotomic polynomial $\Phi_i(q)$ as in Lemma 1. The correspondence between the classes of maximal tori and the F-classes of the Weyl group is well understood for the finite exceptional groups. See [KS02, Shi75, DM87, Sho74, Shi74, DF91, Asc87, FJ93]. For the untwisted groups, the torus orders and corresponding F-class representatives are obtained in Magma [BCP97] (using the command `TwistedTorusOrders`), and the results are obtained by computer. The torus orders for ${}^2E_6(q)$ are obtained from those of $E_6(q)$ by replacing q by $-q$ and the F-class representatives in [FJ93, p.98–99] can be taken to be the same (see also [MT11, Proposition 25.3]). The results for the other twisted groups can be obtained by direct calculation. We provide a detailed example in the case $G = {}^2F_4(q)$ with $q = 2^f$ and f odd.
In this case, there are 11 classes of maximal tori T_i. The orders $|T_i|$, together with the sizes $|C_i|$ of the corresponding F-classes, are given in [Shi75, p. 8]. We list them here for the convenience of the reader:

(1) $|T_1| = (q-1)^2$; $\frac{|C_1|}{|W|} = 1/16$;
(2) $|T_2| = q^2 - 1$; $\frac{|C_2|}{|W|} = 1/4$;
(3) $|T_3| = (q-1)(q - \sqrt{2q} + 1)$; $\frac{|C_3|}{|W|} = 1/8$;
(4) $|T_4| = (q-1)(q + \sqrt{2q} + 1)$; $\frac{|C_4|}{|W|} = 1/8$;
(5) $|T_5| = q^2 + 1$; $\frac{|C_5|}{|W|} = 1/16$;
(6) $|T_6| = (q - \sqrt{2q} + 1)^2$; $\frac{|C_6|}{|W|} = 1/96$;
(7) $|T_7| = (q + \sqrt{2q} + 1)^2$; $\frac{|C_7|}{|W|} = 1/96$;
(8) $|T_8| = (q+1)^2$; $\frac{|C_8|}{|W|} = 1/48$;

e	$\lvert Q\rvert/\lvert{}^2F_4(q)\rvert$
1	$7/16+1/2\phi_{1,r}+1/16\phi_{1,r}^2$
$2(r=3)$	$89/144+1/4\phi_{2,r}+1/48\phi_{2,r}^2$
$2(r>3)$	$35/48+1/4\phi_{2,r}+1/48\phi_{2,r}^2$
4	$77/96+3/16\phi_{4,r}+1/96\phi_{4,r}^2$
6	$5/6+1/6\phi_{6,r}$
12	$11/12+1/12\phi_{12,r}$

Table 2. Proportions of r-regular elements in $G={}^2F_4(q)$ where q has multiplicative order e modulo r ($|Z(G)|=1$)

(9) $|T_9|=q^2-q+1$; $\frac{|C_9|}{|W|}=1/6$;

(10) $|T_{10}|=q^2-\sqrt{2q^3}+q-\sqrt{2q}+1$; $\frac{|C_{10}|}{|W|}=1/12$;

(11) $|T_{11}|=q^2+\sqrt{2q^3}+q+\sqrt{2q}+1$; $\frac{|C_{11}|}{|W|}=1/12$.

If $e=1$, then r divides $q-1=\Phi_1(q)$ and since $r\nmid q$, we have $r\geq 3$. In particular r does not divide $q+1=\Phi_2(q)$. Moreover, we note that

$$(q-\sqrt{2q}+1)(q+\sqrt{2q}+1)=q^2+1=\Phi_4(q)$$

so r does not divide $(q\pm\sqrt{2q}+1)$ either by Lemma 1. Furthermore,

$$(q^2+\sqrt{2q^3}+q+\sqrt{2q}+1)(q^2-\sqrt{2q^3}+q-\sqrt{2q}+1)=q^4-q^2+1=\Phi_{12}(q)$$

so r does not divide $|T_{10}|$ or $|T_{11}|$. Thus we have the proportion $|Q\cap T_i|/|T_i|$ of r-regular elements in T_i is $1/(q-1)^2$ for $i=1$; $1/(q-1)$ for $i=2,3,4$ and 1 for $i=5,\dots,11$. Applying Theorem 2 implies that if $e=1$, then the proportion of r-regular elements in $G={}^2F_4(q)$ is

$$\frac{|Q|}{|G|}=\sum_{i=1}^{11}\frac{|C_i|}{|W|}.\frac{|T_i\cap Q|}{|T_i|}=\frac{1}{16(q-1)_r^2}+\frac{1}{(q-1)_r}\left(\frac{1}{4}+\frac{1}{8}+\frac{1}{8}\right)+\frac{1}{16}+\frac{1}{96}+\frac{1}{96}+\frac{1}{48}+\frac{1}{6}+\frac{1}{12}+\frac{1}{12}$$
$$=\frac{7}{16}+\frac{1}{2(q-1)_r}+\frac{1}{16(q-1)_r^2},$$

which appears in line 1 of Table 2. If $e=2$ then $r\nmid q-1$, $\Phi_{12}(q)$ or $\Phi_4(q)$, we have $(q^2-1)_r=(q+1)_r$ and $(\Phi_6(q))_r=(3,r)$. Thus $|Q\cap T_i|/|T_i|=1$ for $i=1,3,4,5,6,7,10,11$; $1/(q+1)_r$ for $i=2$; $1/(q+1)_r^2$ for $i=8$ and $1/(3,r)$ for $i=9$. Applying Theorem 2 again implies that

$$\frac{|Q|}{|G|}=\sum_{i=1}^{11}\frac{|C_i|}{|W|}.\frac{|T_i\cap Q|}{|T_i|}=\frac{1}{16}+\frac{1}{8}+\frac{1}{8}+\frac{1}{16}+\frac{1}{96}+\frac{1}{96}+\frac{1}{12}+\frac{1}{12}+\frac{1}{4(q+1)_r}+\frac{1}{48(q+1)_r^2}+\frac{1}{6(3,r)}$$
$$=\frac{9}{16}+\frac{1}{6(3,r)}+\frac{1}{4(q+1)_r}+\frac{1}{48(q+1)_r^2},$$

which gives lines 2 and 3 of Table 2. The other cases are similar and we omit the details.

4. Tables for the precise proportions in the simply connected finite exceptional groups

We note that by Lemma 1, the proportion of r-regular elements in the finite *simple* exceptional groups is obtained by multiplying by $|Z(G)|_r$.

e	$\lvert Q\rvert/\lvert {}^2B_2(q)\rvert$
1	$1/2+1/2\phi_{1,r}$
4	$3/4+1/4\phi_{4,r}$

TABLE 3. Proportions of r-regular elements in $G={}^2B_2(q)$ where q has multiplicative order e modulo r ($\lvert Z(G)\rvert=1$)

e	$\lvert Q\rvert/\lvert {}^3D_4(q)\rvert$
1 ($r=2, q\equiv 1 \pmod 4$)	$(17\phi_{1,r}^2+12\phi_{1,r}+4)/48\phi_{1,r}^2$
1 ($r=2, q\equiv 3 \pmod 4$)	$(17\phi_{2,r}^2+12\phi_{2,r}+4)/48\phi_{2,r}^2$
1 ($r=3$)	$(82\phi_{1,r}^2+72\phi_{1,r}+6)/216\phi_{1,r}^2$
1 ($r>3$)	$(10\phi_{1,r}^2+12\phi_{1,r}+2)/24\phi_{1,r}^2$
2 ($r=3$)	$(82\phi_{2,r}^2+72\phi_{2,r}+6)/216\phi_{2,r}^2$
2 ($r>3$)	$(10\phi_{2,r}^2+12\phi_{2,r}+2)/24\phi_{2,r}^2$
3	$(15\phi_{3,r}^2+8\phi_{3,r}+1)/24\phi_{3,r}^2$
6	$(15\phi_{6,r}^2+8\phi_{6,r}+1)/24\phi_{6,r}^2$
12	$(3\phi_{12,r}+1)/4\phi_{12,r}$

TABLE 4. Proportions of r-regular elements in $G={}^3D_4(q)$ where q has multiplicative order e modulo r ($\lvert Z(G)\rvert=1$)

e	$\lvert Q\rvert/\lvert F_4(q)\rvert$
$1(r=2, q\equiv 1 \pmod 4)$	$(3577\phi_{1,r}^4+3696\phi_{1,r}^3+1672\phi_{1,r}^2+192\phi_{1,r}+16)/18432\phi_{1,r}^4$
$1(r=2, q\equiv 3 \pmod 4)$	$(3577\phi_{2,r}^4+3696\phi_{2,r}^3+1672\phi_{2,r}^2+192\phi_{2,r}+16)/18432\phi_{2,r}^4$
$1(r=3)$	$(3337\phi_{1,r}^4+3816\phi_{1,r}^3+1326\phi_{1,r}^2+216\phi_{1,r}+9)/10368\phi_{1,r}^4$
$1(r>3)$	$(385\phi_{1,r}^4+552\phi_{1,r}^3+190\phi_{1,r}^2+24\phi_{1,r}+1)/1152\phi_{1,r}^4$
$2(r=3)$	$(3337\phi_{2,r}^4+3816\phi_{2,r}^3+1326\phi_{2,r}^2+216\phi_{2,r}+9)/10368\phi_{2,r}^4$
$2(r>3)$	$(385\phi_{2,r}^4+552\phi_{2,r}^3+190\phi_{2,r}^2+24\phi_{2,r}+1)/1152\phi_{2,r}^4$
3	$(55\phi_{3,r}^2+16\phi_{3,r}+1)/72\phi_{3,r}^2$
4	$(77\phi_{4,r}^2+18\phi_{4,r}+1)/96\phi_{4,r}^2$
6	$(55\phi_{6,r}^2+16\phi_{6,r}+1)/72\phi_{6,r}^2$
8	$(7\phi_{8,r}+1)/8\phi_{8,r}$
12	$(11\phi_{12,r}+1)/12\phi_{12,r}$

TABLE 5. Proportions of r-regular elements in $G=F_4(q)$ where q has multiplicative order e modulo r ($\lvert Z(G)\rvert=1$)

e	$\lvert Q\rvert/\lvert G_2(q)\rvert$
$1(r=2, q\equiv 1 \pmod 4)$	$(17\phi_{1,r}^2+12\phi_{1,r}+4)/48\phi_{1,r}^2$
$1(r=2, q\equiv 3 \pmod 4)$	$(17\phi_{2,r}^2+12\phi_{2,r}+4)/48\phi_{2,r}^2$
$1(r=3)$	$(9\phi_{1,r}^2+2\phi_{1,r}^2+18\phi_{1,r}+3)/36\phi_{1,r}^2$
$1(r>3)$	$(3\phi_{1,r}^2+2\phi_{1,r}^2+6\phi_{1,r}+1)/12\phi_{1,r}^2$
$2(r=3)$	$(9\phi_{2,r}^2+2\phi_{2,r}^2+18\phi_{2,r}+3)/36\phi_{2,r}^2$
$2(r>3)$	$(3\phi_{2,r}^2+2\phi_{2,r}^2+6\phi_{2,r}+1)/12\phi_{2,r}^2$
3	$(5\phi_{3,r}+1)/6\phi_{3,r}$
6	$(5\phi_{6,r}+1)/6\phi_{6,r}$

TABLE 6. Proportions of r-regular elements in $G=G_2(q)$ where q has multiplicative order e modulo r ($\lvert Z(G)\rvert=1$)

e	$\lvert Q\rvert/\lvert E_6(q)\rvert$
$1\ (r=2, q\equiv 1 \pmod 4)$	$(179840\phi_{1,r}^6+131292\phi_{1,r}^5+113709\phi_{1,r}^4+19080\phi_{1,r}^3+4920\phi_{1,r}^2+288\phi_{1,r}+16)/829440\phi_{1,r}^6$
$1\ (r=2, q\equiv 3 \pmod 4)$	$(17557\phi_{2,r}^4+12024\phi_{2,r}^3+3928\phi_{2,r}^2+288\phi_{2,r}+16)/73728\phi_{2,r}^4$
$1(r=3)$	$(110240\phi_{1,r}^6+489348\phi_{1,r}^5+303003\phi_{1,r}^4+71280\phi_{1,r}^3+9450\phi_{1,r}^2+972\phi_{1,r}+27)/1399680\phi_{1,r}^6$
$1(r=5)$	$(61600\phi_{1,r}^6+90684\phi_{1,r}^5+44709\phi_{1,r}^4+18000\phi_{1,r}^3+2550\phi_{1,r}^2+180\phi_{1,r}+5)/259200\phi_{1,r}^6$
$1(r>5)$	$(12320\phi_{1,r}^6+22284\phi_{1,r}^5+13089\phi_{1,r}^4+3600\phi_{1,r}^3+510\phi_{1,r}^2+36\phi_{1,r}+1)/51840\phi_{1,r}^6$
$2(r=3)$	$(3337\phi_{2,r}^4+3816\phi_{2,r}^3+1326\phi_{2,r}^2+216\phi_{2,r}+9)/10368\phi_{2,r}^4$
$2(r>3)$	$(385\phi_{2,r}^4+552\phi_{2,r}^3+190\phi_{2,r}^2+24\phi_{2,r}+1)/1152\phi_{2,r}^4$
3	$(440\phi_{3,r}^3+183\phi_{3,r}^2+24\phi_{3,r}+1)/648\phi_{3,r}^3$
4	$(77\phi_{4,r}^2+18\phi_{4,r}+1)/96\phi_{4,r}^2$
5	$(4\phi_{5,r}+1)/5\phi_{5,r}$
6	$(55\phi_{6,r}^2+16\phi_{6,r}+1)/72\phi_{6,r}^2$
8	$(7\phi_{8,r}+1)/8\phi_{8,r}$
9	$(8\phi_{9,r}+1)/9\phi_{9,r}$
12	$(11\phi_{12,r}+1)/12\phi_{12,r}$

TABLE 7. Proportions of r-regular elements in $G=E_6(q)$ where q has multiplicative order e modulo r ($\lvert Z(G)\rvert=(3,q-1)$)

e	$\lvert Q\rvert/\lvert{}^2E_6(q)\rvert$
$1, r = 2, q \equiv 1 \pmod 4$	$(17557\phi_{1,r}^4 + 12024\phi_{1,r}^3 + 3928\phi_{1,r}^2 + 288\phi_{1,r} + 16)/73728\phi_{1,r}^4$
$1, r = 2, q \equiv 3 \pmod 4$	$(179840\phi_{2,r}^6 + 131292\phi_{2,r}^5 + 113709\phi_{2,r}^4 + 19080\phi_{2,r}^3 + 4920\phi_{2,r}^2 + 288\phi_{2,r} + 16)/829440\phi_{2,r}^6$
$1(r = 3)$	$(3337\phi_{1,r}^4 + 3816\phi_{1,r}^3 + 1326\phi_{1,r}^2 + 216\phi_{1,r} + 9)/(10368\phi_{1,r}^4)$
$1(r > 3)$	$(385\phi_{1,r}^4 + 552\phi_{1,r}^3 + 190\phi_{1,r}^2 + 24\phi_{1,r} + 1)/1152\phi_{1,r}^4$
$2(r = 3)$	$(110240\phi_{2,r}^6 + 489348\phi_{2,r}^5 + 303003\phi_{2,r}^4 + 71280\phi_{2,r}^3 + 9450\phi_{2,r}^2 + 972\phi_{2,r} + 27)/(1399680\phi_{2,r}^6)$
$2(r = 5)$	$(61600\phi_{2,r}^6 + 90684\phi_{2,r}^5 + 44709\phi_{2,r}^4 + 18000\phi_{2,r}^3 + 2550\phi_{2,r}^2 + 180\phi_{2,r} + 5)/259200\phi_{2,r}^6$
$2(r > 5)$	$(12320\phi_{2,r}^6 + 22284\phi_{2,r}^5 + 13089\phi_{2,r}^4 + 3600\phi_{2,r}^3 + 510\phi_{2,r}^2 + 36\phi_{2,r} + 1)/51840\phi_{2,r}^6$
3	$(55\phi_{3,r}^2 + 16\phi_{3,r} + 1)/72\phi_{3,r}^2$
4	$(77\phi_{4,r}^2 + 18\phi_{4,r} + 1)/96\phi_{4,r}^2$
6	$(440\phi_{6,r}^3 + 183\phi_{6,r}^2 + 24\phi_{6,r} + 1)/648\phi_{6,r}^3$
8	$(7\phi_{8,r} + 1)/8\phi_{8,r}$
10	$(4\phi_{10,r} + 1)/5\phi_{10,r}$
12	$(11\phi_{12,r} + 1)/12\phi_{12,r}$
18	$(8\phi_{18,r} + 1)/9\phi_{18,r}$

TABLE 8. Proportions of r-regular elements in $G = {}^2E_6(q)$ where q has multiplicative order e modulo r ($\lvert Z(G)\rvert = (3, q+1)$)

e	$\lvert Q\rvert/\lvert E_7(q)\rvert$
$1\ (r=2, q\equiv 1 \pmod 4)$	$(41419665\phi_{1,r}^7+95510014\phi_{1,r}^6+30219588\phi_{1,r}^5+10204152\phi_{1,r}^4+952560\phi_{1,r}^3+116256\phi_{1,r}^2+4032\phi_{1,r}+128)/371589120\phi_{1,r}^7$
$1\ (r=2, q\equiv 3 \pmod 4)$	$(41419665\phi_{2,r}^7+95510014\phi_{2,r}^6+30219588\phi_{2,r}^5+10204152\phi_{2,r}^4+952560\phi_{2,r}^3+116256\phi_{2,r}^2+4032\phi_{2,r}+128)/371589120\phi_{2,r}^7$
$1(r=3)$	$(20191815\phi_{1,r}^7+23513057\phi_{1,r}^6+12786039\phi_{1,r}^5+3532473\phi_{1,r}^4+405405\phi_{1,r}^3+31563\phi_{1,r}^2+1701\phi_{1,r}+27)/78382080\phi_{1,r}^7$
$1(r=5)$	$(3828825\phi_{1,r}^7+6047743\phi_{1,r}^6+2739177\phi_{1,r}^5+621159\phi_{1,r}^4+108675\phi_{1,r}^3+8085\phi_{1,r}^2+315\phi_{1,r}+5)/14515200\phi_{1,r}^7$
$1(r=7)$	$(5360355\phi_{1,r}^7+7764581\phi_{1,r}^6+4647699\phi_{1,r}^5+1140573\phi_{1,r}^4+152145\phi_{1,r}^3+11319\phi_{1,r}^2+441\phi_{1,r}+7)/20321280\phi_{1,r}^7$
$1(r>7)$	$(765765\phi_{1,r}^7+1286963\phi_{1,r}^6+663957\phi_{1,r}^5+162939\phi_{1,r}^4+21735\phi_{1,r}^3+1617\phi_{1,r}^2+63\phi_{1,r}+1)/2903040\phi_{1,r}^7$
$2(r=3)$	$(20191815\phi_{2,r}^7+23513057\phi_{2,r}^6+12786039\phi_{2,r}^5+3532473\phi_{2,r}^4+405405\phi_{2,r}^3+31563\phi_{2,r}^2+1701\phi_{2,r}+27)/78382080\phi_{2,r}^7$
$2(r=5)$	$(3828825\phi_{2,r}^7+6047743\phi_{2,r}^6+2739177\phi_{2,r}^5+621159\phi_{2,r}^4+108675\phi_{2,r}^3+8085\phi_{2,r}^2+315\phi_{2,r}+5)/14515200\phi_{2,r}^7$
$2(r=7)$	$(5360355\phi_{2,r}^7+7764581\phi_{2,r}^6+4647699\phi_{2,r}^5+1140573\phi_{2,r}^4+152145\phi_{2,r}^3+11319\phi_{2,r}^2+441\phi_{2,r}+7)/20321280\phi_{2,r}^7$
$2(r>7)$	$(765765\phi_{2,r}^7+1286963\phi_{2,r}^6+663957\phi_{2,r}^5+162939\phi_{2,r}^4+21735\phi_{2,r}^3+1617\phi_{2,r}^2+63\phi_{2,r}+1)/2903040\phi_{2,r}^7$
3	$(935\phi_{3,r}^3+327\phi_{3,r}^2+33\phi_{3,r}+1)/1296\phi_{3,r}^3$
4	$(77\phi_{4,r}^2+18\phi_{4,r}+1)/96\phi_{4,r}^2$
5	$(9\phi_{5,r}+1)/10\phi_{5,r}$
6	$(935\phi_{6,r}^3+327\phi_{6,r}^2+33\phi_{6,r}+1)/1296\phi_{6,r}^3$
7	$(13\phi_{7,r}+1)/14\phi_{7,r}$
8	$(7\phi_{8,r}+1)/8\phi_{8,r}$
9	$(17\phi_{9,r}+1)/18\phi_{9,r}$
10	$(9\phi_{10,r}+1)/10\phi_{10,r}$
12	$(11\phi_{12,r}+1)/12\phi_{12,r}$
14	$(13\phi_{14,r}+1)/14\phi_{14,r}$
18	$(17\phi_{18,r}+1)/18\phi_{18,r}$

TABLE 9. Proportions of r-regular elements in $G=E_7(q)$ where q has multiplicative order e modulo r ($\lvert Z(G)\rvert=(2,q-1)$)

e	$\lvert Q\rvert/\lvert E_8(q)\rvert$
$1\ (r=2, q\equiv 1 \pmod 4)$	$(41492380321\phi_{1,r}^8 + 27525566640\phi_{1,r}^7 + 16480551440\phi_{1,r}^6 + 2132907840\phi_{1,r}^5 + 295921248\phi_{1,r}^4 + 14434560\phi_{1,r}^3 + 815360\phi_{1,r}^2 + 15360\phi_{1,r} + 256)/178362777600\phi_{1,r}^8$
$1\ (r=2, q\equiv 3 \pmod 4)$	$(41492380321\phi_{2,r}^8 + 27525566640\phi_{2,r}^7 + 16480551440\phi_{2,r}^6 + 2132907840\phi_{2,r}^5 + 295921248\phi_{2,r}^4 + 14434560\phi_{2,r}^3 + 815360\phi_{2,r}^2 + 15360\phi_{2,r} + 256)/178362777600\phi_{2,r}^8$
$1(r=3)$	$(16277566921\phi_{1,r}^8 + 21789381960\phi_{1,r}^7 + 7567769940\phi_{1,r}^6 + 1361503080\phi_{1,r}^5 + 159928398\phi_{1,r}^4 + 8913240\phi_{1,r}^3 + 366660\phi_{1,r}^2 + 9720\phi_{1,r} + 81)/56435097600\phi_{1,r}^8$
$1(r=5)$	$(5363541841\phi_{1,r}^8 + 7507077000\phi_{1,r}^7 + 2845718900\phi_{1,r}^6 + 501215400\phi_{1,r}^5 + 53801790\phi_{1,r}^4 + 4095000\phi_{1,r}^3 + 150500\phi_{1,r}^2 + 3000\phi_{1,r} + 25)/17418240000\phi_{1,r}^8$
$1(r=7)$	$(1509595087\phi_{1,r}^8 + 2115252600\phi_{1,r}^7 + 761301260\phi_{1,r}^6 + 172854360\phi_{1,r}^5 + 18315906\phi_{1,r}^4 + 1146600\phi_{1,r}^3 + 42140\phi_{1,r}^2 + 840\phi_{1,r} + 7)/4877107200\phi_{1,r}^8$
$1(r>7)$	$(215656441\phi_{1,r}^8 + 323507400\phi_{1,r}^7 + 130085780\phi_{1,r}^6 + 24693480\phi_{1,r}^5 + 2616558\phi_{1,r}^4 + 163800\phi_{1,r}^3 + 6020\phi_{1,r}^2 + 120\phi_{1,r} + 1)/696729600\phi_{1,r}^8$
$2(r=3)$	$(16277566921\phi_{2,r}^8 + 21789381960\phi_{2,r}^7 + 7567769940\phi_{2,r}^6 + 1361503080\phi_{2,r}^5 + 159928398\phi_{2,r}^4 + 8913240\phi_{2,r}^3 + 366660\phi_{2,r}^2 + 9720\phi_{2,r} + 81)/56435097600\phi_{2,r}^8$
$2(r=5)$	$(5363541841\phi_{2,r}^8 + 7507077000\phi_{2,r}^7 + 2845718900\phi_{2,r}^6 + 501215400\phi_{2,r}^5 + 53801790\phi_{2,r}^4 + 4095000\phi_{2,r}^3 + 150500\phi_{2,r}^2 + 3000\phi_{2,r} + 25)/17418240000\phi_{2,r}^8$
$2(r=7)$	$(1509595087\phi_{2,r}^8 + 2115252600\phi_{2,r}^7 + 761301260\phi_{2,r}^6 + 172854360\phi_{2,r}^5 + 18315906\phi_{2,r}^4 + 1146600\phi_{2,r}^3 + 42140\phi_{2,r}^2 + 840\phi_{2,r} + 7)/4877107200\phi_{2,r}^8$
$2(r>7)$	$(215656441\phi_{2,r}^8 + 323507400\phi_{2,r}^7 + 130085780\phi_{2,r}^6 + 24693480\phi_{2,r}^5 + 2616558\phi_{2,r}^4 + 163800\phi_{2,r}^3 + 6020\phi_{2,r}^2 + 120\phi_{2,r} + 1)/696729600\phi_{2,r}^8$
3	$(124729\phi_{3,r}^4 + 28400\phi_{3,r}^3 + 2310\phi_{3,r}^2 + 80\phi_{3,r} + 1)/155520\phi_{3,r}^4$
4	$(31345\phi_{4,r}^4(5,r) + 2304\phi_{4,r}^4 + 11100\phi_{4,r}^3(5,r) + 1270\phi_{4,r}^2(5,r) + 60\phi_{4,r}(5,r) + (5,r))/46080\phi_{4,r}^4(5,r)$
5	$(551\phi_{5,r}^2 + 48\phi_{5,r} + 1)/600\phi_{5,r}^2$
6	$(124729\phi_{6,r}^4 + 28400\phi_{6,r}^3 + 2310\phi_{6,r}^2 + 80\phi_{6,r} + 1)/155520\phi_{6,r}^4$
7	$(13\phi_{7,r} + 1)/14\phi_{7,r}$
8	$(161\phi_{8,r}^2 + 30\phi_{8,r} + 1)/192\phi_{8,r}^2$
9	$(17\phi_{9,r} + 1)/18\phi_{9,r}$
10	$(551\phi_{10,r}^2 + 48\phi_{10,r} + 1)/600\phi_{10,r}^2$
12	$(253\phi_{12,r}^2 + 34\phi_{12,r} + 1)/288\phi_{12,r}^2$
14	$(13\phi_{14,r} + 1)/14\phi_{14,r}$
15	$(29\phi_{15,r} + 1)/30\phi_{15,r}$
18	$(17\phi_{18,r} + 1)/18\phi_{18,r}$
20	$(19\phi_{20,r} + 1)/20\phi_{20,r}$
24	$(23\phi_{24,r} + 1)/24\phi_{24,r}$
30	$(29\phi_{30,r} + 1)/30\phi_{30,r}$

TABLE 10. Proportions of r-regular elements in $G = E_8(q)$ where q has multiplicative order e modulo r ($\lvert Z(G)\rvert = 1$)

e	$\lvert Q\rvert/\lvert {}^2G_2(q)\rvert$
$1\ (r=2)$	$7/12+1/6(q+1)_2$
$1\ (r>3)$	$1/2+1/2\phi_{1,r}$
2	$5/6+1/6\phi_{2,r}$
6	$5/6+1/6\phi_{6,r}$

Table 11. Proportions of r-regular elements in $G = {}^2G_2(q)$ where q has multiplicative order e modulo r ($|Z(G)| = 1$)

References

[Asc87] Michael Aschbacher, *Chevalley groups of type G_2 as the group of a trilinear form*, J. Algebra **109** (1987), no. 1, 193–259. MR 898346 (88g:20089)

[BCP97] Wieb Bosma, John Cannon, and Catherine Playoust, *The* Magma *algebra system I: The user language*, J. Symbolic Comput. **24** (1997), no. 3-4, 235–265.

[BGPW] László Babai, Simon Guest, Cheryl E. Praeger, and Robert A. Wilson, *Proportions of r-regular elements in finite classical groups*, J. London Math. Soc., to appear.

[BPS09] László Babai, Péter P. Pálfy, and Jan Saxl, *On the number of p-regular elements in finite simple groups*, LMS J. Comput. Math. **12** (2009), 82–119.

[Car85] Roger W. Carter, *Finite groups of Lie type*, Pure and Applied Mathematics (New York), John Wiley & Sons Inc., New York, 1985. MR 794307 (87d:20060)

[DF91] D. I. Deriziotis and A. P. Fakiolas, *The maximal tori in the finite Chevalley groups of type E_6, E_7 and E_8*, Comm. Algebra **19** (1991), no. 3, 889–903. MR 1102992 (92c:20081)

[DM87] D. I. Deriziotis and G. O. Michler, *Character table and blocks of finite simple triality groups ${}^3D_4(q)$*, Trans. Amer. Math. Soc. **303** (1987), no. 1, 39–70. MR 896007 (88j:20011)

[FJ93] Peter Fleischmann and Ingo Janiszczak, *The semisimple conjugacy classes of finite groups of Lie type E_6 and E_7*, Comm. Algebra **21** (1993), no. 1, 93–161. MR 1194553 (93k:20029)

[IKS95] I. M. Isaacs, William M. Kantor, and N. Spaltenstein, *On the probability that a group element is p-singular*, Journal of Algebra **176** (1995), no. 1, 139–181.

[KS02] William M. Kantor and Ákos Seress, *Prime power graphs for groups of Lie type*, J. Algebra **247** (2002), no. 2, 370–434. MR 1877859 (2003a:20026)

[Leh98] G. I. Lehrer, *The cohomology of the regular semisimple variety*, J. Algebra **199** (1998), no. 2, 666–689. MR 1489931 (98k:20080)

[MT11] Gunter Malle and Donna Testerman, *Linear algebraic groups and finite groups of Lie type*, Cambridge Studies in Advanced Mathematics, vol. 133, Cambridge University Press, Cambridge, 2011. MR 2850737 (2012i:20058)

[NP10] Alice C. Niemeyer and Cheryl E. Praeger, *Estimating proportions of elements in finite groups of Lie type*, Journal of Algebra **324** (2010), no. 1, 122 – 145.

[Shi74] Kenichi Shinoda, *The conjugacy classes of Chevalley groups of type F_4 over finite fields of characteristic* 2, 133–159.

[Shi75] ———, *The conjugacy classes of the finite Ree groups of type F_4*, 1–15.

[Sho74] Toshiaki Shoji, *The conjugacy classes of Chevalley groups of type F_4 over finite fields of characteristic $p \neq 2$*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **21** (1974), 1–17.

School of Mathematics, University of Southampton, Southampton SO17 1BJ, UK
E-mail address: `s.d.guest@soton.ac.uk`